Lincomycin 10%

Composition:

Each ml solution contains: 100 mg lincomycin (as HCl)

Action:

Lincomycin is a small spectrum bacteriostatic antibiotic, active against Grampositive bacteria, anaerobes and mycoplasmata.

Indications:

Pigs: Dysentery, Mycoplasma infections (infectious arthritis, pneumonia), E.coli enteritis, Erysipelas.

Calves: Enteritis and pneumonia.

Sheep and goats: Dermatitis contagiosa interdigitalis ovis (footrot).

Poultry: CRD, E. coli tracheitis, enteritis (Salmonella, Treponema, E. coli), arthritis,

septicemia, sepsis, omphalitis.

Administration/dosage:

By intramuscular or subcutaneous (poultry) injection:

Pigs, calves, sheep: 1 ml per 10 kg b.w. daily for 3-7 days

Cats and dogs: 0.5 - 1 ml per 5 kg b.w. once or twice daily for 3-5 days

Poultry: 0.5 ml per 2.5 kg b.w.. s.c.

Contra-Indications:

Animals hypersensitive to antibiotics. Do not administer to rabbits.

Advised withdrawal times:

Meat: 7 days.

Conservation:

Cool (8 - 15°C), protected from light.

Presentation:

Per 12 vials of 100 ml.